

National Library  
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	MIM
Search PubMed	for					Go	Clear
Limits		Preview/Index		History		Clipboard	
Display	Abstract	Save	Text	Order	Add to Clipboard		

RS1 565

Entrez PubMed

☐ 1: *J Pharm Pharmacol* 1998 Jun;50(6):567-74[Related Articles, Books, LinkOut](#)

PubMed Services

## Use of transgenic animals in understanding molecular mechanisms of toxicity.

**Wolf CR, Henderson CJ**

Imperial Cancer Research Fund, Molecular Pharmacology Unit, Biomedical Research Centre, Ninewells Hospital and Medical School, Dundee, UK.

Related Resources

Understanding molecular mechanisms of chemical toxicity and the potential risks of drugs to man is a pivotal part of the drug development process. With the dramatic increase in the number of new chemical entities arising from high throughput screening, there is an urgent need to develop systems for the rapid evaluation of potential drugs so that those agents which are most likely to be free of adverse effects can be identified at the earliest possible stage in drug development. The complex mechanisms of action of chemical toxins has made it extremely difficult to evaluate the precise toxic mechanism and also the relative role of specific genes in either potentiating or ameliorating the toxic effect. This problem can be addressed by the application of genetic strategies. Such strategies can exploit strain differences in susceptibility to specific toxic agents or, with the rapidly developing technologies, can exploit the use of transgenic animals where specific genes can be manipulated and subsequent effects on chemical toxicity evaluated. Transgenic animals can be exploited in a variety of ways to understand mechanisms of chemical toxicity. For example, a human gene encoding a drug metabolizing enzyme can be directly introduced and the effects on toxic response evaluated. Alternatively, specific genes can be deleted from the mouse genome and the consequences on toxicological response determined. Many toxic chemical agents modulate patterns of gene expression within target cells. This can be used to screen for responses to different types of toxic insult. In such experiments the promoter of a stress-regulated gene can be ligated to a suitable reporter gene, such as lacZ, or green fluorescent protein, and inserted into the genome of an appropriate test species. On administration of a chemical agent, cells which are sensitive to the toxic effects of that chemical will express the reporter, which can then be identified using an appropriate assay system. This latter strategy provides the potential for screening a large number of compounds rapidly for their potential toxic effects and also provides information on tissue and cellular specificity. Experiments using transgenic animals can be complex, and care must be taken to ensure that the results are not affected by background activities within the species being used. For example, the introduction of a specific human cytochrome P450 gene may have no effect on the metabolic disposition of a drug or toxin because of the

background activity within the mouse. As the toxicity of a chemical agent is determined by a wide range of different factors including drug uptake, metabolism, detoxification and repair, differences between man and the species being used could potentially generate a toxic response in the animal model whereas no toxicity may be observed in man. In spite of these confounding factors, the application of transgenic animals to toxicological issues has enormous potential for speeding up the drug discovery process and will undoubtedly become part of this process in the future.

Publication Types:

- Review
- Review, tutorial

PMID: 9680065, UI: 98343659

Display	Abstract	<input type="checkbox"/>	Save	Text	Order	Add to Clipboard
---------	----------	--------------------------	------	------	-------	------------------

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)

=> d his

(FILE 'HOME' ENTERED AT 16:16:07 ON 26 JAN 2001)

FILE 'MEDLINE' ENTERED AT 16:16:15 ON 26 JAN 2001

```

L1      0 S (TRANSGENIC MICE OR TRANSGENIC ANIMALS OR TRANSGENIC MOUSE) (
L2      0 S (TRANSGENIC MICE OR TRANSGENIC ANIMALS OR TRANSGENIC MOUSE) A
L3      175 S (TRANSGENIC MICE OR TRANSGENIC ANIMALS OR TRANSGENIC MOUSE) A
L4      124 S (TRANSGENIC MICE OR TRANSGENIC ANIMALS OR TRANSGENIC MOUSE) A
L5      99 S (TRANSGENIC MICE OR TRANSGENIC ANIMALS OR TRANSGENIC MOUSE) (
L6      262861 S REVIEW
L7      0 S L5 AND L6
L8      0 S (TRANSGENIC MICE OR TRANSGENIC ANIMALS OR TRANSGENIC MOUSE) (
L9      1439 S (HEAT SHOCK OR STRESS INDUC?) (S) PROMOTER
L10     0 S L7 AND L9
L11     0 S L5 AND L9
L12     1 S L5 AND STRESS
L13     1 S LUCIFERAS ASSAY AND IN VIVO
L14     3 S L5 AND LUCIFERASE ASSAY

```

=> s heat shock or stress induc?

```

106050 HEAT
363 HEATS
106258 HEAT
      (HEAT OR HEATS)
78298 SHOCK
3315 SHOCKS
80102 SHOCK
      (SHOCK OR SHOCKS)
16241 HEAT SHOCK
      (HEAT(W)SHOCK)
168411 STRESS
10214 STRESSES
173724 STRESS
      (STRESS OR STRESSES)
1182142 INDUC?
8301 STRESS INDUC?
      (STRESS(W)INDUC?)
L15     23838 HEAT SHOCK OR STRESS INDUC?

```

=> s 15 and 115

```

L16     1 L5 AND L15

```

=> d ibib abs